

AD-A093 574

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER

F/G 12/1

ELEMENTARY PROOFS OF AN INEQUALITY FOR SYMMETRIC FUNCTIONS FOR --ETC(U)

AUG 80 R ZIELKE

DAA629-80-C-0041

UNCLASSIFIED

MRC-TSR-2113

NL

1 2 1
4 6 3 5 7 4

END

DATA
FILMED

DTIC

AD A093574

MRC Technical Summary Report #2113

ELEMENTARY PROOFS OF AN INEQUALITY
FOR SYMMETRIC FUNCTIONS FOR $n \leq 5$

Roland Zielke

(1)
p. 5

LEVEL

Mathematics Research Center
University of Wisconsin-Madison
610 Walnut Street
Madison, Wisconsin 53706

August 1980

(Received June 24, 1980)

DIS
SELECTED
JAN 7 1981
C

Approved for public release
Distribution unlimited

DDC FILE COPY

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

80 12 22 057

(14) 11-2-T-113

11-2-T-113

UNIVERSITY OF WISCONSIN - MADISON
MATHEMATICS RESEARCH CENTER

(15) 2

ELEMENTARY PROOFS OF AN INEQUALITY
FOR SYMMETRIC FUNCTIONS FOR $n \leq 5$.

10 Roland Zielke

+

val to

9 Technical Summary Report #2113
August 1980

ABSTRACT

(15) 11-2-T-113

For $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ let the elementary symmetric functions

$\psi_j = \mathbb{R}^n \rightarrow \mathbb{R}$ be defined by

$$\psi_j(x) = \sum_{1 \leq i_1 < \dots < i_j \leq n} x_{i_1} \dots x_{i_j}, \quad j = 1, \dots, n. \text{ So the real polynomial } p_x \text{ of}$$

degree n with leading coefficient 1 and zeros in $-x_1, \dots, -x_n$ is given by $p_x(t) = t^n + \sum_{i=1}^n \psi_i(x) t^{n-i}$.

Let $x, y \in \mathbb{R}_+^n$ be points with $\psi_i(x) \leq \psi_i(y)$ for $i = 1, \dots, n$.

It was conjectured (see [2]) that this implies $\psi_i(x^\alpha) \leq \psi_i(y^\alpha)$

for every $\alpha \in (0, 1]$ and $i = 1, \dots, n$, where x^α is defined by

$$x^\alpha = (x_1^\alpha, \dots, x_n^\alpha).$$

By an argument involving total positivity, this conjecture may be reduced to the problem of finding a piecewise differentiable path

$\{\phi(t) | t \in [0, 1]\}$ in \mathbb{R}_+^n with $\phi(0) = x$, $\phi(1) = y$ and such that

$\psi_i(\phi(t))$ is monotone increasing with t for each $i = 1, \dots, n$ (see [1]).

This problem looks deceptively simple but was only recently solved

by Efroymson, Swartz and Wendroff using a rather involved argument.

We give elementary proofs for $n \leq 5$.

AMS (MOS) Subject Classification: 26D05

Key Words: real polynomials, inequalities

Work Unit Number 3 (Numerical Analysis and Computer Science)

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041 and in part by a travel grant of the Deutsche Forschungsgemeinschaft.

SIGNIFICANCE AND EXPLANATION

Some aspects of the heat transfer in the emergency cooling of nuclear reactors lead to a nonlinear eigenvalue problem, the so-called model quench front problem. Laquer and Wendroff suggested a procedure for computing bounds of the eigenvalue which depend - among other things - on the validity of a certain inequality for elementary symmetric functions. This inequality is of interest in itself and was recently proved by Efroymson, Swartz and Wendroff using a fairly complicated argument. We give an elementary proof for $n \leq 5$.

Accession For	<input checked="" type="checkbox"/>
NTIS GRA&I	<input type="checkbox"/>
ERIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist. and/or	
Point	Special
A	

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

ELEMENTARY PROOFS OF AN INEQUALITY
FOR SYMMETRIC FUNCTIONS FOR $n \leq 5$

Roland Zielke

Let $\mathbb{R}_{+}^n = \{z \in \mathbb{R}^n \mid \bigwedge_i z_i \geq 0\}$ and $\Delta_{+}^n = \{z \in \mathbb{R}_{+}^n \mid z_1 \leq z_2 \leq \dots \leq z_n\}$.

Let $\sigma : \mathbb{R}^n \rightarrow \mathbb{R}^n$, $\sigma : x \rightarrow \sigma(x)$, be defined by

$$\prod_{i=1}^n (t - x_i) = t^n + \sum_{i=1}^n \sigma_i(x) t^{n-i} =: p_x(t) \text{ for } t \in \mathbb{R}.$$

So we have $\sigma_i(x) = \sum_{1 \leq j_1 < \dots < j_i \leq n} (-1)^i x_{j_1} \dots x_{j_i}$, $i = 1, \dots, n$,

$$\text{and } \sigma(\mathbb{R}_{+}^n) \subset \mathbb{R}_{+}^n.$$

Let \mathbb{R}^n be partially ordered by " $x < y$ iff $x_i \leq y_i$ for $i = 1, \dots, n$

and $x \neq y$ ". Let $x, y \in \Delta_{+}^n$ be points with $\sigma(x) < \sigma(y)$ and

$M = \{z \in \Delta_{+}^n \mid \sigma(x) \leq \sigma(z) \leq \sigma(y)\}$. So M is compact.

Theorem A: a) There is a continuous mapping $\phi : [0, 1] \rightarrow M$ with

$\phi(0) = x$, $\phi(1) = y$ and $\sigma(\phi(u)) < \sigma(\phi(v))$ for all $u, v \in [0, 1]$ with $u < v$.

b) ϕ is continuously differentiable except on a finite set.

By an argument involving total positivity (see [1]) one may derive from theorem A the following result:

Theorem B: If z^α is defined by $z^\alpha = (-|z_1|^\alpha, \dots, -|z_n|^\alpha)$ for $z \in \Delta_{+}^n$ and $\alpha \in \mathbb{R}$, we have $\sigma(x^\beta) \leq \sigma(y^\beta)$ for $\beta \in (0, 1]$.

Subsequently we shall prove theorem A for $n \leq 5$.

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041 and in part by a travel grant of the Deutsche Forschungsgemeinschaft.

Proof: a) It is sufficient to find a $\delta > 0$ and a $g \in \mathbb{P}_{n-1}$ with nonnegative coefficients such that $p_x + \lambda g$ has n nonpositive real zeros $x_1^{(\lambda)}, \dots, x_n^{(\lambda)}$ for all $\lambda \in [0, \delta]$ and $\sigma(x^{(\lambda)})$ is strictly increasing for $\lambda \in [0, \delta]$.

For $x_n < x_{n-1} < \dots < x_1$ the claim is trivial. Also trivial is the following

Lemma 1: If $y < x$ and $y_1 < 0$, then $\sigma_i(x) < \sigma_i(y)$ for all i .

We denote $d := p_y - p_x$. We consider the cases $n = 2, 3, 4, 5$ separately:

$n = 2$:

$x_2 = x_1$: choose $g(t) = t$, if $\sigma_1(x) < \sigma_1(y)$.

If $\sigma_1(x) = \sigma_1(y)$, we have $d(t) = \alpha$ for some $\alpha > 0$, and p_y has no zeros, a contradiction.

$n = 3$:

case 1: $x_3 < x_2 = x_1$: choose $g(t) = t$, if $\sigma_1(x) < \sigma_1(y)$.

Otherwise we have $d(t) = \alpha + \beta t^2$ for some $\alpha, \beta \in \mathbb{R}_+$, and lemma 1 gives a contradiction.

case 2: $x_3 = x_2 < x_1$: choose $g(t) = 1$, if $\sigma_0(x) < \sigma_0(y)$;

choose $g(t) = t^2$ if $\sigma_2(x) < \sigma_2(y)$.

Otherwise we have $d(t) = \alpha t$ for some $\alpha \in \mathbb{R}_+$, implying $x_i < y_i$ for $i = 1, 2, 3$, a contradiction.

case 3: $x_1 = x_2 = x_3$: choose $g(t) = t(t - x_1)$, if $\sigma_i(x) < \sigma_i(y)$ for $i = 1, 2$.

Otherwise, if $\sigma_1(x) = \sigma_1(y)$, go to $n=3$, case 1.

If $\sigma_2(x) = \sigma_2(y)$, consider p'_x, p'_y and go to $n=2$.

$n = 4$:

case 1: $x_4 < x_3 = x_2 < x_1$: choose $g(t) = 1$, if $\sigma_0(x) < \sigma_0(y)$;

choose $g(t) = t^2$, if $\sigma_2(x) < \sigma_2(y)$.

Otherwise we have $d(t) = \alpha t + \beta t^3$ for some $\alpha, \beta \in \mathbb{R}_+$.

$\Rightarrow d'(t) = \alpha + 3\beta t^2 \Rightarrow p'_y(t) > p'_x(t)$ and $p'_y(t) > p'_x(t)$ for $t \in (-\infty, 0)$.

So all zeros of p'_y are smaller than all zeros of p'_x , yielding $\sigma_2(x) < \sigma_2(y)$, a contradiction.

case 2: a) $x_4 < x_3 < x_2 = x_1$ or b) $x_4 = x_3 < x_2 < x_1$:

choose $g(t) = t$, if $\sigma_1(x) < \sigma_1(y)$;

choose $g(t) = t^3$, if $\sigma_3(x) < \sigma_3(y)$;

otherwise we have $d(t) = \alpha + \beta t^2$, so $d > 0$, $d' < 0$, $d'' > 0$ on $(-\infty, 0)$ and $d'(0) = 0$.

For a) this implies $Z(p'_y) \subset (-\infty, x_3)$, but also $Z(p'_y) \cap (x_1, 0) \neq \emptyset$, a contradiction.

For b) this implies that either all zeros of p'_y are larger than all zeros of p'_x , or that all zeros of p''_y are larger than all zeros of p''_x , in both cases a contradiction.

case 3: $x_4 < x_3 = x_2 = x_1$: choose $g(t) = t(t-x_1)$, if $\sigma_1(x) < \sigma_1(y)$ and $\sigma_2(x) < \sigma_2(y)$.

Otherwise, if $\sigma_2(x) = \sigma_2(y)$, we have $d(t) = \alpha + \beta t + \gamma t^3$, so $d'(t) = \beta + 3\gamma t^2$. Now go to n=3, case 1.

If $\sigma_1(x) = \sigma_1(y)$, we have $d(t) = \alpha + \beta t^2 + \gamma t^3$. So d has only one zero z in $(-\infty, 0)$, d' has only one zero z' in $(-\infty, 0)$, $d'(0) = 0$, $z \leq z' \leq 0$.

If p_1 has no zero in $(x_1, 0)$, the same holds for p'_y . But then all zeros of p'_y are smaller than all zeros of p'_x , and lemma 1 gives a contradiction.

If p_y has a zero in $(x_1, 0)$ we have $z \in (x_1, 0)$ and thus $p_x \cdot p_y$ and $p'_x \cdot p'_y$ on $(-\infty, z)$. But then again the zeros of p'_y are smaller than those of p'_x .

case 4: $x_4 = x_3 = x_2 = x_1$: choose $g(t) = t - x_2$, if $\sigma_0(x) < \sigma_0(y)$ and $\sigma_1(x) < \sigma_1(y)$; choose $g(t) = t^2(t - x_2)$, if $\sigma_2(x) < \sigma_2(y)$ and $\sigma_3(x) < \sigma_3(y)$.

Otherwise: a) If $\sigma_0(x) = \sigma_0(y)$ and $\sigma_2(x) = \sigma_2(y)$, go to $n = 4$, case 1.

b) If $\sigma_1(x) = \sigma_1(y)$ and $\sigma_3(x) = \sigma_3(y)$, go to $n = 4$, case 2b.

c) If $\sigma_0(x) = \sigma_0(y)$ and $\sigma_3(x) = \sigma_3(y)$, we have $d(t) = \alpha t + \beta t^2$ and $\alpha, \beta > 0$ w.l.o.g.. So p_1'' has its zeros in (x_2, x_1) , and d has its negative zero in $(x_1, 0)$. But then $x_i \leq y_i$ for all i , a contradiction.

d) If $\sigma_1(x) = \sigma_1(y)$ and $\sigma_2(x) = \sigma_2(y)$, we have $d(t) = \alpha + \beta t^3$ and $\alpha, \beta > 0$ w.l.o.g. If d had its zero z in $(-\infty, x_1]$, we would have

$y_i = x_i$ for all i in contradiction to lemma 1. $\Rightarrow z \in (x_1, 0)$. Let

z_1 be the local minimum of p_x . We have $p_y' > 0$ in $[z_1, 0]$, so

$Z(p_y') \subset (-\infty, x_2)$, for otherwise p_y would have two local extrema in (x_2, z) with no zero in between. But this again yields a contradiction to lemma 1.

case 5: $x_4 = x_3 = x_2 = x_1$: choose $g(t) = t(t - x_1)^2$, if $\sigma_i(x) < \sigma_i(y)$ for $i = 1, 2, 3$.

Otherwise, if $\sigma_i(x) = \sigma_i(y)$ for $i = 2$ or $i = 3$, consider p_x' and p_y' , i.e., go to $n=3$, case 3.

If $\sigma_1(x) = \sigma_1(y)$, we have $d(t) = \alpha + \beta t^2 + \gamma t^3$ with $\beta, \gamma > 0$ w.l.o.g.

Go to $n=4$, case 3, corresponding case.

$n = 5$:

We use the following notations:

The zeros of p_x' are z_4, z_3, z_2, z_1 with $z_4 \leq z_3 \leq z_2 \leq z_1$.

The zeros of p_x'' are w_3, w_2, w_1 , with $w_3 \leq w_2 \leq w_1$.

The negative zeros of d are p, q, r, \dots with $p \leq q \leq r \leq \dots$

The negative zeros of d' are p', q', r' with $p' \leq q' \leq r'$

The negative zeros of d'' are p'', q'' with $p'' \leq q''$.

The statement " $\sigma_i(x) = \sigma_i(y)$ " is called A_i , $i = 0, 1, \dots, 4$.

α, β, γ are nonnegative real numbers.

case 1:

a) $x_5 < x_4 < x_3 < x_2 = x_1$

b) $x_5 < x_4 = x_3 < x_2 < x_1$

c) $x_5 < x_4 = x_3 < x_2 = x_1$

Choose $g(t) = t$ if A_1 , choose $g(t) = t^3$ if A_3 .

If $\neg A_1 \wedge \neg A_3$, we have $d(t) = \alpha + \beta t^2 + \gamma t^4 \Rightarrow d'(0) = 0 \wedge d' < 0 < d''$ on $(-\infty, 0)$.

a) We have $Z(p_Y) \subset (-\infty, x_5) \cup (x_4, x_3)$ and $Z(p'_Y) \cap (x_1, 0) \neq \emptyset \Rightarrow Z(p_Y) \cap (x_1, 0) \neq \emptyset$, contradiction.

c) Follows from a).

b) We have $Z(p_Y) \subset (-\infty, x_5) \cup (x_2, x_1)$ and $Z(p'_Y) \subset (-\infty, z_4) \cup (x_3, z_2) \cup (z_1, 0)$.

If $Z(p''_Y) \subset (-\infty, w_3]$, lemma 1 gives a contradiction to $\neg A_3$.

$$\Rightarrow \#(Z(p''_Y) \cap (-\infty, w_3]) = 1 \wedge \#(Z(p''_Y) \cap [w_2, w_1]) = 2$$

$\Rightarrow p'_Y$ has 2 zeros in (z_3, z_1)

$\Rightarrow p'_Y$ has 2 zeros in (z_3, z_2)

$\Rightarrow p_Y$ has 1 zeros in $(z_3, z_2) \subset (x_3, x_2)$, contradiction.

case 2:

a) $x_5 < x_4 < x_3 = x_2 < x_1$

b) $x_5 = x_4 < x_3 < x_2 < x_1$

c) $x_5 = x_4 < x_3 = x_2 < x_1$

Choose $g(t) = 1$ if A_0 ,

$g(t) = t^2$ if A_2 ,

$g(t) = t^4$ if A_4 .

If $\neg A_0 \wedge \neg A_2 \wedge \neg A_4$, we have $d(t) = \alpha t + \beta t^3 \Rightarrow d' > 0 > d''$ on $(-\infty, 0)$.

a) p_Y has one zero in $(x_1, 0)$ and 4 zeros in $(x_5, x_4) \wedge Z(p'_Y) \subset (z_4, z_3)$

$\Rightarrow Z(p''_Y) \subset (z_4, z_3)$. But $p''_Y(0) = p''_X(0) \geq 0 \wedge p''_Y(w_1) < p''_X(w_1) = 0 \Rightarrow Z(p''_Y) \cap (w_1, 0) \neq \emptyset$, contradiction.

b) p_Y has one zero in $(x_1, 0)$ and 4 zeros in $(x_3, x_2) \wedge Z(p'_Y) \subset (z_2, z_1)$

$\Rightarrow Z(p''_Y) \subset (w_1, z_1)$, contradiction.

case 3:

a) $x_5 \leq x_4 < x_3 = x_2 = x_1$

b) $x_5 = x_4 = x_3 < x_2 \leq x_1$

a) Choose $g(t) = t(t-x_1)$ if $A_1 \wedge A_2$,

$$g(t) = t^3(t-x_1) \text{ if } A_3 \wedge A_4,$$

$$g(t) = t(t^3-x_1^3) \text{ if } A_1 \wedge A_4. \text{ Otherwise we have:}$$

a) 1) $\neg A_2 \wedge \neg A_4$: consider p'_x, p'_y and go to n=4, case 2a).

a) 2) $\neg A_1 \wedge \neg A_3$: we have $d(t) = \alpha + \beta t^2 + \gamma t^4$ and $d' < 0$ on $(-\infty, 0) = Z(p'_y) \cap (x_1, 0) \neq \emptyset$. But $Z(p'_y) \subset (-\infty, x_1) \Rightarrow Z(p'_y) \subset (-\infty, x_1)$ contradiction.

a) 3) $\neg A_1 \wedge \neg A_4$: we have $d(t) = \alpha + \beta t^2 + \gamma t^3$ and $\gamma > 0$ w.l.o.g. So d, d', d'' have exactly one negative zero each, and $p < p' < p''$.

$$\left. \begin{array}{l} \text{If } p \in [x_1, 0] \Rightarrow Z(p_y) \subset (x_5, x_4) \cup (x_1, p) \Rightarrow Z(p'_y) \subset (z_4, z_3) \\ \text{If } p < x_1 \Rightarrow Z(p_y) \subset (-\infty, x_1) \Rightarrow Z(p'_y) \subset (-\infty, x_1) \Rightarrow \\ p' \in (x_1, 0) \Rightarrow Z(p'_y) \subset (z_4, z_3) \end{array} \right\} \Rightarrow$$

$Z(p''_y) \subset (z_4, z_3)$. But $p'' \in (x_1, 0) \Rightarrow p''_y(x_1) < 0 \Rightarrow Z(p''_y) \cap (x_1, 0) \neq \emptyset$, contradiction.

b) Choose $g(t) = t(t-x_5)$ if $A_1 \wedge A_2$,

$$g(t) = t^3(t-x_5) \text{ if } A_3 \wedge A_4,$$

$$g(t) = t(t^3-x_5^3) \text{ if } A_1 \wedge A_4. \text{ Otherwise we have:}$$

b) 1) $\neg A_2 \wedge \neg A_4$: consider p'_x, p'_y and go to n=4, case 2b).

b) 2) $\neg A_1 \wedge \neg A_3$: we have $d(t) = \alpha + \beta t^2 + \delta t^4$, so $d' < 0 < d''$ on $(-\infty, 0)$ and

$$Z(p_y) \subset (-\infty, x_5) \cup (x_2, x_1),$$

$$Z(p'_y) \subset (-\infty, z_2) \cup (z_1, 0),$$

$$Z(p''_y) \subset (-\infty, x_5) \cup (w_2, w_1).$$

p''_y has one zero in $(-\infty, x_5)$ and two zeros in (w_2, w_1) , for otherwise lemma 1 and A_3 give a contradiction.

b) 2) a) $Z(p_y) \subset (-\infty, x_5) \Rightarrow$ lemma 1 contradicts $\neg A_3$.

b) 2) b) $\#(Z(p_Y) \cap (x_2, x_1)) = 2 \Rightarrow \#(Z(p'_Y) \cap (z_1, 0)) = 1$
 $\Rightarrow p'_1$ has 3 zeros in $(-\infty, x_5)$
 $\Rightarrow p''_Y$ has 2 zeros in $(-\infty, x_5)$, contradiction.

b) 2) c) $\#(Z(p_Y) \cap (x_2, x_1)) = 4 = \#(Z(p'_Y) \cap (z_1, x_1)) = 3$
 $\Rightarrow \#(Z(p''_Y) \cap (z_1, x_1)) = 2$, contradiction

b) 3) $\neg A_1 \wedge \neg A_4$: we have $d(t) = \alpha + \beta t^2 + \gamma t^3$.

b) 3) a) $p' \in (z_1, 0)$: If $\#(Z(p'_Y) \cap (z_1, 0)) = 2$, lemma 1 and $\neg A_4$ give a contradiction.

If $Z(p'_Y) \subset (-\infty, z_1) \Rightarrow Z(p'_Y) \subset (z_2, z_1]$. From $d'' < 0$ in $(-\infty, p')$ follows $Z(p''_Y) \subset (w_1, z_Y)$, contradiction.

b) 3) b) $p' \in [z_2, z_1) \Rightarrow$ lemma 1 and $\neg A_4$ give a contradiction.

b) 3) c) $p' \in (x_5, z_2) \Rightarrow Z(p'_Y) \subset (p', z_2) \cup (z_1, 0)$
 $\Rightarrow \#(Z(p'_Y) \cap (p', z_2)) = 3 \Rightarrow \#(Z(p_Y) \cap (p', z_2)) = 2$.

But $p_Y > p_X > 0$ in $(p, z_2) \cap (x_5, z_2) \supset (p', z_2)$, contradiction.

b) 3) d) $p' < x_5$: If $\#(Z(p'_Y) \cap (x_5, z_2)) = 2 \Rightarrow Z(p_Y) \cap (x_5, z_2) \neq \emptyset$,
but $p_Y > p_X > 0$ on (x_5, z_2) , contradiction.

case 4:

$x_5 < x_4 = x_3 = x_2 < x_1$: Choose $g(t) = t - x_3$ if $A_0 \wedge A_1$,
 $g(t) = t^2(t - x_3)$ if $A_2 \wedge A_3$,
 $g(t) = t^3 - x_3^3$ if $A_0 \wedge A_3$. Otherwise

we have:

1) $\neg A_1 \wedge \neg A_3$: consider p'_X, p'_Y and go to n=4, case 1.

2) $\neg A_0 \wedge \neg A_2$: we have $d(t) = \alpha t + \beta t^3 + \gamma t^4$.

$\Rightarrow d, d', d''$, have each exactly one negative zero, and $p < p' < p''$.

One checks that $Z(p''_Y) \subset (x_3, 0)$ or $Z(p''_Y) \subset (-\infty, x_3]$ are impossible.

a) $p' \in (z_1, 0) \Rightarrow Z(p'_Y) \subset (-\infty, z_4) \cup (z_1, 0)$.

If p'_Y had 3 zeros in (z_1, p') , p''_Y would have 2 zeros in (z_1, p') , contradiction.

If p'_Y had 3 zeros in $(-\infty, z_4)$, p''_Y would have 2 zeros in $(-\infty, z_4)$

$\Rightarrow Z(p''_Y) \subset (-\infty, w_3) \Rightarrow \neg A_2$ and lemma 1 give a contradiction.

b) $p' \in (x_3, z_1)$:

If p'_Y has exactly one zero $> p'$, p''_Y has at most one zero $\geq x_3$.

So p''_Y has no zero $\geq x_3$, contradiction.

If p'_Y has 3 zeros in $(p', 0)$, p_Y has 4 zeros in $(x_1, 0)$.

$\Rightarrow Z(p_Y) \subset (x_1, 0)$, contradiction.

c) $p' \in (-\infty, x_3) \Rightarrow \#(Z(p'_Y) \cap (x_3, z_1)) \geq 2$, for otherwise

$Z(p''_Y) \subset (-\infty, x_3)$, contradiction. So we have $\#(Z(p_Y) \cap (x_3, z_1)) \geq 1$.

but this contradicts $p_Y < p_X < 0$ in (x_3, z_1) .

3) $\neg A_0 \wedge A_3$: we have: $d(t) = \alpha t + \beta t^2 + \gamma t^4$. Then d and d' have exactly one negative zero each, $p < p'$, and $d'' > 0$ on $(-\infty, 0]$ w.l.o.g..

So p''_Y has 2 zeros in (x_3, w_1) , i.e., p'_Y has a local maximum r and a local minimum s with $p' < r < s < w_1$.

$\Rightarrow p_Y$ has a local maximum $l \in (r, s)$, and $d(l) > 0$.
For $\varepsilon > 0$ sufficiently small, $d(-\varepsilon) < 0$ $\} \Rightarrow p \in (l, 0) \Rightarrow p' < p$, contradiction.

case 5:

$x_5 = x_4 < x_3 < x_2 = x_1$: choose $g(t) = t(t-x_3)$ if $A_1 \wedge A_2$,
 $g(t) = t^3(t-x_3)$ if $A_3 \wedge A_4$,
 $g(t) = t(t^3-x_3)$ if $A_1 \wedge A_4$. Other-

wise we have:

1) $\neg A_1 \wedge \neg A_3$: go to n=5, case 1a.

2) $\neg A_2 \wedge \neg A_4$: we have $d(t) = \alpha + \beta t + \gamma t^3$, so $d' > 0 > d''$ on $(-\infty, 0)$.

$\Rightarrow Z(p'_Y) \subset (x_5, z_3) \cup (z_2, x_1) \wedge Z(p''_Y) \subset (w_3, w_2) \cup (w_1, 0)$

$\Rightarrow p'_Y$ has at least 2 zeros in (z_2, x_1) and 1 zero in (x_5, w_3)

$\Rightarrow p_Y$ has a local minimum in (z_2, x_1) and a local maximum in (x_5, w_3) .

$\Rightarrow d$ has at least 2 zeros in $(-\infty, 0)$, contradiction.

3) $\neg A_1 \wedge \neg A_4$: we have $d(t) = \alpha + \beta t^2 + \gamma t^3$.

$\Rightarrow d, d', d''$ have each exactly one negative zero, and $p < p' < p''$.

We have either $Z(p_Y) \subset (-\infty, x_3]$ or $Z(p_Y) \subset (x_3, 0)$:

$Z(p_Y) \subset (-\infty, x_3]$ implies $Z(p_Y'') \subset (-\infty, x_3) \Rightarrow p' > x_1 \Rightarrow p'' > x_1$ contradiction.

$Z(p_Y) \subset (x_3, 0)$ implies: $Z(p_Y) \subset (z_2, 0) \Rightarrow Z(p_Y'') \subset (z_2, 0) \Rightarrow Z(p_Y'') \subset (z_2, w_1)$

$\Rightarrow p'' < w_1, \Rightarrow p' < w_1 \Rightarrow Z(p_Y') \cap (x_1, 0] \neq \emptyset \Rightarrow x_1 < p < p' < w_1$ contradiction.

case 6:

$x_5 = x_4 = x_3 = x_2 < x_1$: choose $g(t) = (t-x_5)^2$ if $A_0 \wedge A_1 \wedge A_2$
 $g(t) = t^2(t-x_5)^2$ if $A_2 \wedge A_3 \wedge A_4$
 $g(t) = (t^2+x_5t+x_5^2)(t-x_5)^2 =$
 $= t^4 - x_5t^3 - x_5^3t + x_5^4$, if $A_0 \wedge A_1 \wedge A_3 \wedge A_4$.

Otherwise we have:

1) $(\neg A_1 \wedge \neg A_3)$ or $(\neg A_1 \wedge \neg A_4)$: consider p'_x, p'_y and go to n=4, case 4.

2) $\neg A_0 \wedge \neg A_3$: go to n=5, case 4,3.

3) $\neg A_0 \wedge \neg A_4$: we have $d(t) = \alpha t + \beta t^2 + \gamma t^3$.

$\Rightarrow d$ has 2, d' has 2, d'' has 1 negative zero, and $p < p' < q < q' < 0, p' < p'' < q'$.

From $p'' < w_1$ follows that p_Y'' has at least 2 zeros in $(\max\{p'_1, x_5\}, w_1)$.

So p_Y' has a local maximum r and a local minimum s with $p'' < r$ and

$x_5 < r < s < w_1. \Rightarrow d'(r) > 0 \Rightarrow$ either $r' < q' < r$ or $r < p' < q'$.

As p_Y has a local maximum $l \in (r, s)$ and $d(l) > 0$, we have $l \in (p, q)$,

so $r < l < q < q'$, and so finally $r < p' < q' \Rightarrow r < p''$, contradiction.

case 7:

$x_5 < x_4 = x_3 = x_2 = x_1$: choose $g(t) = t(t-x_1)^2$ if $A_1 \wedge A_2 \wedge A_3$.

Otherwise we have:

1) $\neg A_2$ or $\neg A_3$: consider p'_x, p'_y and go to n=4, case 3.

2) $\neg A_1$: we have $d(t) = \alpha + \beta t^2 + \gamma t^3 + \delta t^4$ with $\delta, \gamma \neq 0$ w.l.o.g..

$\Rightarrow d$ has at most 2 zeros in $(-\infty, 0)$. If d had no zero or one double zero in $(-\infty, 0)$, lemma 1 and $\neg A_1$ would give a contradiction.

$\Rightarrow d$ has exactly 2 zeros in $(-\infty, 0)$, as well as d' and d'' , and $p < p' < q < q' < 0$ and $p' < p'' < q' < q'' < 0$.

claim 1: $Z(p_Y'') \subset (-\infty, w_3] \Rightarrow d''$ has no zero in $[x_1, 0]$.

Proof: explicit computation gives $p_X'' < p_Y''$ on $[x_1, 0]$.

claim 2: $x_5 \leq q \Rightarrow Z(p_Y') \subset (-\infty, q')$.

claim 3: $p' \leq x_1 \leq q' \Rightarrow$ either $p' < z_4$, or p_Y' has 2 zeros in $(q', 0)$.

Proof: If p_Y' has less than 2 zeros in $(q', 0)$, p_Y' has no zero there, so $Z(p_Y') \subset (-\infty, x_1]$. If now p_Y' had a zero $\leq z_4$, lemma 1 and $\neg A_1$ would yield a contradiction.

From $q \leq x_5$ would follow A_1 by lemma 1, a contradiction. So we have $x_5 \leq q \Rightarrow p' < x_1$, for otherwise $x_1 < p' < q' \Rightarrow Z(p_Y') \subset (x_1, p')$ because of claim 2 $\Rightarrow Z(p_Y'') \subset (x_1, p')$, contradiction.

a) $x_1 < q \Rightarrow x_1 < q' \Rightarrow Z(p_Y') \subset (-\infty, x_1)$
b) $q \leq x_1 \Rightarrow Z(p_Y') \subset (-\infty, x_1) \Rightarrow Z(p_Y') \subset (-\infty, x_1)$ } $\Rightarrow Z(p_Y') \subset (z_4, x_1)$, for
otherwise lemma 1 and $\neg A_1$
yield a contradiction.

$\Rightarrow Z(p_Y'') \subset (-\infty, x_1) \wedge x_1 < q' < q'' \Rightarrow x_1 < p'' \Rightarrow Z(p_Y'') \subset (-\infty, w_3)$ contradiction to claim 1.

case 8:

$x_5 = x_4 = x_3 = x_2 = x_1$: choose $g(t) = t(t-x_1)^3$ if $A_1 \wedge A_2 \wedge A_3 \wedge A_4$.

1) $\neg A_2$ or $\neg A_3$ or $\neg A_4$: p_X' and p_Y' can be treated as n=4, case 5.

2) $\neg A_1$: same as n=5, case 7.

b) Let $\| \cdot \|$ denote any fixed norm in \mathbb{R}^n .

We construct $f_1, f_2, \dots, \in P_n$ with corresponding zeros $x^{(1)}, x^{(2)}, \dots, x^{(n)}$ as follows: Let $f_0 = p_X$ and $x^{(0)} = x$. If for $k \geq 0$, $x^{(k)}$ and f_k are

given, for every

$$g \in S := \{f \in \mathbb{P}_{n-1} \mid \max_{t \in [0,1]} |f(t)| = 1\},$$

let δ_g be maximal such that

a) for all $\lambda \in [0, \delta_g]$, $f_k + \lambda g$ has n zeros $z_{g_1}^{(\lambda)}, \dots, z_{g_n}^{(\lambda)}$

with $z_g^{(\lambda)} \in M$

b) $\sigma(z_g^{(\lambda)})$ is strictly increasing for $\lambda \in [0, \delta_g]$.

Let $\hat{g} \in S$ be a function with

$$\|z_{\hat{g}}^{(\delta_{\hat{g}})} - \sigma(f_k)\| = \max_{g \in S} \|z_g^{(\delta_g)} - \sigma(f_k)\|,$$

and define $f_{k+1} = f_k + \delta_{\hat{g}} \hat{g}$, $x^{(k+1)} = z_{\hat{g}}^{(\delta_{\hat{g}})}$.

So p_x and every f_k are connected by a path along which σ is strictly increasing, and this path corresponds to a polygonal arc in $\sigma(M)$ with corners $\sigma(x^{(0)}), \sigma(x^{(1)}), \dots, \sigma(x^{(k)})$. We have to show $f_k = p_y$ occurs for some k .

Suppose the contrary, i.e. $\sigma(x^{(k)}) < \sigma(y)$ for all $k = 1, 2, \dots$.

As $\{\sigma(x^{(k)})\}$ is an increasing sequence, $\sigma^\infty := \lim_{k \rightarrow \infty} \sigma(x^{(k)})$ exists.

Let $x^\infty := \lim_{k \rightarrow \infty} x^{(k)}$, so $\sigma^\infty = \sigma(x^\infty)$, and f_∞ the corresponding polynomial.

There is a $g \in \mathbb{P}_{n-1}$, and a $\delta > 0$ such that $f_\infty + \lambda g$ has n zeros $z_1^{(\lambda)}, \dots, z_n^{(\lambda)}$ with $z^{(\lambda)} \in \Delta_-^n$ for all $\lambda \in [0, 2\delta]$, and $\sigma(z^{(\lambda)})$ is strictly increasing with $\lambda \in [0, 2\delta]$. Let $\alpha = \|\sigma(z^{(\delta)}) - \sigma(x^{(\infty)})\|$.

We shall show that for every $\epsilon > 0$, there is an index n and a

$\tilde{g} \in \mathbb{P}_{n-1}$ (near g) such that

1) $f_k + \lambda \tilde{g}$ has n zeros $\tilde{z}_1^{(\lambda)}, \dots, \tilde{z}_n^{(\lambda)}$ with $\tilde{z}^{(\lambda)} \in \Delta_-^n$ for all $\lambda \in [0, \delta]$,

2) $\sigma(\tilde{z}^{(\lambda)})$ is strictly increasing for $\lambda \in [0, \delta]$,

$$3) \quad \| \sigma(z^{(\delta)}) - \sigma(\tilde{z}^{(\delta)}) \| < \varepsilon.$$

(This implies $\| \sigma(x^{(k+1)}) \| \geq \| \sigma(\tilde{z}^{(\delta)}) \| \geq \| \sigma^\infty \| + \alpha - \varepsilon > \| \sigma^\infty \|$

for all sufficiently small $\varepsilon > 0$, a contradiction.)

Let $\tilde{\varepsilon} > 0$ be arbitrarily fixed and k so large that

$$|(f_\infty - f_k)(t)| < \tilde{\varepsilon} \text{ for all } t \in I := [2x_n^\infty - 1, 1], \text{ and}$$

$$\| x^{(k)} - x^\infty \| < \tilde{\varepsilon}.$$

So in an $\tilde{\varepsilon}$ -neighbourhood of every zero z of f^∞ of multiplicity m , f_k has exactly m zeros counting multiplicities.

As the functions g in part a) of the proof were constructed only in view of the multiplicities of the zeros of f^∞ , \tilde{g} can be constructed correspondingly in view of the zeros of f_k .

As an example, we consider the case $n=5$, case 8 (leaving the analogous details of the other cases to the reader):

For $f_\infty(t) = (t - x_1^\infty)^5$, we had $g(t) = (t - x_1)^3 t$.

For $f_k(t) = \prod_{i=1}^5 (t - x_i^{(k)})$ with $x_5^{(k)} \leq x_4^{(k)} \leq \dots \leq x_1^{(k)}$, we choose

$$\tilde{g}(t) = (t - x_2^{(k)}) (t - x_3^{(k)}) (t - x_4^{(k)}) t$$

$$\Rightarrow \max_{t \in I} \{ |(g - \tilde{g})(t)| \} = O(\tilde{\varepsilon}), \text{ and}$$

$$\max_{t \in I} \{ |(f_\infty + \delta g)(t) - (f_k + \delta \tilde{g})(t)| \} = O(\tilde{\varepsilon}).$$

As $f_\infty + \delta g$ has 2 simple zeros $\neq x_1$, $f_k + \delta \tilde{g}$ has simple zeros near these.

For sufficiently small $\tilde{\varepsilon}$ and large k , statement 3) above holds, too.

References

- [1] Efroymsen, G.A., Swartz, B. and B. Wendroff: An inequality for symmetric functions. Submitted to "Advances in Mathematics".
- [2] Laquer, H.T. and B. Wendroff: Bounds for the model quench front; to appear.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 2113	2. GOVT ACCESSION NO. AD-A093	3. RECIPIENT'S CATALOG NUMBER 574
4. TITLE (and Subtitle) ELEMENTARY PROOFS OF AN INEQUALITY FOR SYMMETRIC FUNCTIONS FOR $n \leq 5$		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
7. AUTHOR(s) Roland Zielke		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53706		8. CONTRACT OR GRANT NUMBER(s) DAA029-80-C-0041
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Research Triangle Park, North Carolina 27709		10. PROGRAM ELEMENT PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 3 - Numerical Analysis and Computer Science
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE August 1980
		13. NUMBER OF PAGES 13
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) real polynomials, inequalities		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) For $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ let the elementary symmetric functions $\psi_j = \mathbb{R}^n \rightarrow \mathbb{R}$ be defined by $\psi_j(x) = \sum_{1 \leq i_1 < \dots < i_j \leq n} x_{i_1} \dots x_{i_j}, \quad j = 1, \dots, n.$ So the real polynomial p_x of degree n (continued)		

ABSTRACT (cont.)

with leading coefficient 1 and zeros in $-x_1, \dots, -x_n$ is given by

$$p_x(t) = t^n + \sum_{i=1}^n \psi_i(x) t^{n-i}.$$

Let $x, y \in \mathbb{R}_+^n$ be points with $\psi_i(x) \leq \psi_i(y)$ for $i = 1, \dots, n$.

It was conjectured (see [2]) that this implies $\psi_i(x^\alpha) \leq \psi_i(y^\alpha)$ for every $\alpha \in (0, 1]$ and $i = 1, \dots, n$, where x^α is defined by $x^\alpha = (x_1^\alpha, \dots, x_n^\alpha)$.

By an argument involving total positivity, this conjecture may be reduced to the problem of finding a piecewise differentiable path $\{\phi(t) | t \in [0, 1]\}$ in \mathbb{R}_+^n with $\phi(0) = x$, $\phi(1) = y$ and such that $\psi_i(\phi(t))$ is monotone increasing with t for each $i = 1, \dots, n$ (see [1]). This problem looks deceptively simple but was only recently solved by Efroymson, Swartz and Wendroff using a rather involved argument. We give elementary proofs for $n \leq 5$.